

READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

COOPERS® PERMATRACE® COPPER 10



CAPSULES FOR CALVES AND ADULT CATTLE

ACTIVE CONSTITUENT: 390 g/kg COPPER OXIDE (CuO)
Sustained release copper capsules for prevention and treatment of copper deficiency
in cattle over 3 months of age. Each capsule contains 10 g sustained release copper particles.

100 CAPSULES

FLA51 1905/00

COOPERS®
EST. 1843

COOPERS® PERMATRACE® COPPER 10

CAPSULES FOR CALVES AND ADULT CATTLE

DIRECTIONS FOR USE

Contraindications:

Not to be administered to cattle of unknown copper status or those treated with other sources of copper.
Not to be used in conjunction with selenium supplementation unless deficiency of both copper and selenium has been confirmed.
This product is contraindicated for use in cattle suffering from jaundice and other liver disorders such as liver fluke, or stock grazing plants containing toxic alkaloids, eg: Heliotrope, Ragwort, Patersons Curse (Salvation Jane).
This product is contraindicated for use in calves less than 3 months of age (i.e. when rumen is developed).

Seek advice of veterinary or cattle husbandry specialist to ascertain copper deficiency and type of treatment required for cattle under consideration.

DOSAGE:

Important: Do not use excessive force when dosing in order to avoid trauma and damage to the throat of the animal.

Calves, up to 200 kg (under 1 year old) – one capsule per head annually.
Adult cattle, over 200 kg – 2 capsules per head or 1 capsule of COOPERS PERMATRACE COPPER 20 annually.

WITHHOLDING PERIOD: Zero (0) days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days.

GENERAL INSTRUCTIONS:

Unless corrective treatment is required immediately, the preferred time of treatment is late Winter or early Spring.
To be effective the capsule must be swallowed. It is recommended that capsules be administered using the Coopers® Cattle Pellet Applicator. It is important to carefully read the applicator use instruction leaflet enclosed in container before use. In areas where liver fluke are endemic stock should be treated with a highly effective flukicide at least 3 weeks before dosing with this copper supplement.

For more information, call Coopers Customer Service on 1 800 226 511.

SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with the eyes and skin. Wash hands after use.
FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SPECIALIST ADVICE IN EMERGENCY ONLY

Coopers Animal Health (Intervet Australia Pty Limited)

1 800 226 511

ALL HOURS – AUSTRALIA-WIDE

Warranty: Intervet Australia Pty Limited (IAPL), also known as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

DISPOSE of empty container by wrapping with paper and putting in garbage.

STORE below 30°C (Room Temperature). Store in original container in a safe place. Keep out of reach of children.

BATCH

EXPIRY

APVMA Approval No.: 47689/55234

COOPERS ANIMAL HEALTH

(Intervet Australia Pty Limited)

91-105 Harpin Street, Bendigo East, VIC 3550

® Registered Trademark

® Yellow (PANTONE® 123) the predominant colour of

Coopers packaging is a Registered Trademark.

PANTONE is a Registered Trademark of Pantone, Inc.

© 2013

100 CAPSULES



FLA51 1905/00