


173861 R3

**PLUS SELENIUM FOR LAMBS**

**TASVAX 5 in 1**  
COOPERS®  
**VACCINE FOR SHEEP, CATTLE AND GOATS**

9 1316147 930751

**POISON**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY



**COOPERS®**  
**TASVAX 5 in 1**  
**VACCINE FOR SHEEP, CATTLE AND GOATS**

**PLUS SELENIUM FOR LAMBS**

**500 mL**

**ACTIVE CONSTITUENTS:**  
*Clostridium perfringens* Type D - toxoid  
*Clostridium chauvoei* - toxoid and inactivated cells  
*Clostridium septicum* - toxoid  
*Clostridium novyi* Type B - toxoid and inactivated cells  
*Clostridium tetani* - toxoid  
 0.5 g/L SELENIUM as SODIUM SELENATE  
 Contains 0.15 mg/mL THIOMERSAL as preservative.

**Potency Units**  
 5 U/mL  
 Meets Ph.Eur.  
 2.5 U/mL  
 3.5 U/mL  
 2.5 U/mL

CLOSTRIDIAL VACCINE for immunity against Pulpy Kidney, Blackleg, Malignant Oedema, Black Disease and Tetanus in sheep, cattle and goats and to protect against selenium deficiency in lambs.

**COOPERS®**   
EST. 1843

**COOPERS®**  
**TASVAX 5 in 1**  
**VACCINE FOR SHEEP, CATTLE AND GOATS**

**PLUS SELENIUM FOR LAMBS**

**For immunity against**

- Pulpy Kidney (*Clostridium perfringens* Type D)
- Blackleg (*Clostridium chauvoei*)
- Malignant Oedema (*Clostridium septicum*)
- Black Disease (*Clostridium novyi* Type B)
- Tetanus (*Clostridium tetani*)

in sheep, cattle and goats and to protect against selenium deficiency in lambs.

**SHEEP, LAMBS and GOATS: 250 DOSES**  
**CATTLE: 125 DOSES**

109, 500, 101  
109, 500, 101 x 50, 500, 101 x 2, 490, 101  
MSD, 101, 101 + 101, 101 + 101, 101 + 101

**READ THE ENCLOSED LEAFLET FOR FULL INSTRUCTIONS.**  
**DIRECTIONS FOR USE**  
**Contraindications:** This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.  
**Dosage and administration**  
**Shake contents before use. Use all product within 24 hours of opening.** Do not expose directly to heat and light during use. The use of an insulated container is recommended during the day to protect the vaccine. If it is to be hung beside the race, ensure it is on the shady side.  
**Dose:** Sheep, lambs and goats: 2 mL by subcutaneous injection.  
 Cattle: 4 mL by subcutaneous injection.  
 Inject only under the skin and not into the muscle, by raising a fold of skin and injecting the dose under the raised skin - **NOT INTO IT.** Cattle, sheep and goats should be injected high on the side of the neck, behind the ear.  
**General directions**  
**CAUTION: AVOID CARCASS DAMAGE:**  
 1. Sterilise all injection apparatus by boiling for at least ten minutes before use. Avoid use of strong disinfectants on apparatus.  
 2. Maintain maximum cleanliness at all times.  
 3. Keep needles sharp and clean. Change needles frequently (every 20-30 animals). Use short needles. Recommended maximum sizes 15 mm x 18 gauge for sheep and goats; 15 mm x 16 gauge for cattle.  
 4. Check that the equipment is functioning properly and that it is delivering the correct dose.  
 5. As far as possible avoid injection of animals in wet weather or under dusty conditions.  
 6. Inject only under the skin and not into the muscle, by raising a fold of skin and injecting the dose under the raised skin - **NOT INTO IT.**  
 7. Cattle, sheep and goats should be injected high on the side of the neck, behind the ear.  
**WITHHOLDING PERIOD: Zero (0) days.**  
**TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days.**

**FIRST AID**  
 If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.  
 If skin contact occurs, remove contaminated clothing and wash skin thoroughly.  
**ADDITIONAL USER SAFETY INFORMATION**  
 Take care to avoid self-injection. In the event of self-administration, seek medical attention if you are concerned. Show the package leaflet or label to the Medical Practitioner.  
 Additional information is listed in the safety data sheet.  
**DISPOSAL:** Dispose of empty container by wrapping with paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container.  
**STORAGE:** Store between 2°C and 8°C (refrigerate, do not freeze). Protect from light.  
 APVMA Approval No.: 61165/109609  
 Ⓟ **EXP. REFER TO BATCH AND EXPIRY ON TOP PANEL**  
**COOPERS ANIMAL HEALTH**  
 Intervet Australia Pty Limited  
 (trading as MSD Animal Health)  
 91-105 Harpin Street, Bendigo East VIC 3550  
 Phone: 1800 226 511  
 www.coopersanimalhealth.com.au  
 Ⓟ Registered Trademark.  
 Ⓟ Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.

**COOPERS®**  
**TASVAX 5 in 1**  
**VACCINE FOR SHEEP, CATTLE AND GOATS**

**PLUS SELENIUM FOR LAMBS**



**500 mL**

173861 R3

**PLUS SELENIUM FOR LAMBS**

**TASVAX 5 in 1**  
COOPERS®  
**VACCINE FOR SHEEP, CATTLE AND GOATS**

9 1316147 930751