

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY


COOPERS[®]
PANACUR[®] 100

ORAL ANTHELMINTIC FOR CATTLE AND HORSES



ACTIVE CONSTITUENT:
100 g/L FENBENDAZOLE

Ready to use broad spectrum worm drench for control of benzimidazole sensitive mature and immature roundworms and lungworm in cattle, and for the control of large strongyles, small strongyles, ascarids and pinworms in horses.



F-LA3001124/02

COOPERS

5 Litres

COOPERS® PANACUR® 100

ORAL ANTHELMINTIC FOR CATTLE AND HORSES

COOPERS PANACUR 100 contains fenbendazole, a member of the benzimidazole family of drenches. COOPERS PANACUR 100 is active against both mature (adult) and immature (larval) fenbendazole sensitive nematode parasites. COOPERS PANACUR 100 has activity (80-90%) against inhibited (hypobiotic) larvae of *Ostertagia* in cattle.

COOPERS PANACUR 100 is effective against sensitive strains of the following internal parasites:

Cattle:

Barbers Pole Worm (*Haemonchus* spp.), Small Brown Stomach Worm (*Ostertagia* spp.), Stomach Hair Worm (*Trichostrongylus* spp.), Small Intestinal Worm (*Cooperia* spp.), Thin Necked Intestinal Worm (*Nematodirus* spp.), Nodule Worm (*Oesophagostomum radiatum*), Whipworm (*Trichuris* spp.), *Capillaria* spp., Lungworms (*Dictyocaulus viviparus*), Hookworm (*Bunostomum phlebotomum*).

COOPERS PANACUR 100 also aids in the control of Tapeworm (*Moniezia* spp.)

Horses:

Large Strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*), Small Strongyles (*Cyathostomum* spp., *Tridontophorus* spp.), Ascarids (*Parascaris equorum*), Pinworms (*Oxyuris equi*).

COOPERS PANACUR 100 does not stain or predispose animals to photosensitisation. COOPERS PANACUR 100 is palatable and readily accepted by livestock. COOPERS PANACUR 100 may be administered concurrently with vaccines, pour-on lice treatments and dipping operations.

Resistance may develop to any chemical.

DIRECTIONS FOR USE

Shake container before use: COOPERS PANACUR 100 is ready to use, no dilution or mixing is necessary. COOPERS PANACUR 100 may be administered orally through standard drenching equipment. After use, wash equipment in clean water.

Recommendations for use

Cattle

Dose Rate Cattle: 3.75mL COOPERS PANACUR 100 per 50kg bodyweight (7.5mg fenbendazole/kg bodyweight).

Dose Rate Guide Cattle:

Liveweight kg	Dose Volume (mL)	Number of doses per 5L Pack	Liveweight kg	Dose Volume (mL)	Number of doses per 5L Pack
Up to 50	3.75	1333	351-400	30.0	166
51-100	7.5	667	401-450	33.75	148
101-150	11.25	444	451-500	37.5	133
151-200	15.0	333	501-550	41.25	121
201-250	18.75	267	551-600	45.0	111
251-300	22.5	222	601-650	48.75	102
301-350	26.25	190			

For animals over 650kg dose at rate of 3.75mL per 50kg bodyweight.

A representative sample of animals should be weighed before treatment either with scales or weighband. Dose according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers etc.) Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

For the control of Liver Fluke (*Fasciola hepatica*) in cattle as well as the above mentioned internal parasites. COOPERS PANACUR 100 can be mixed with Fasinex® or Fascicare® - see following table.

The mixture is stable for up to 12 months at 30°C (room temperature). Shake mixture vigorously for 2 minutes or use mechanical mixing prior to use. **This mixture should not be used in horses.**

COOPERS PANACUR 100 Mixed with		
Fasinex® 100	Fascicare® 120	Fasinex® 240
Mixing Instructions		
Mix 5 parts COOPERS PANACUR 100 with 8 parts Fasinex® 100	Mix 3 parts COOPERS PANACUR 100 with 4 parts Fascicare® 120	Mix 3 parts COOPERS PANACUR 100 with 2 parts Fasinex® 240
COOPERS PANACUR 100 at 3.75mL per 50kg plus Fasinex® 100 at 6mL per 50kg bodyweight.	COOPERS PANACUR 100 at 3.75mL per 50kg plus Fascicare® 120 at 5mL per 50kg bodyweight.	COOPERS PANACUR 100 at 3.75mL per 50kg plus Fasinex® 240 at 2.5mL per 50kg bodyweight.
Mixture should be given at 9.75mL per 50kg bodyweight.	Mixture should be given at 8.75mL per 50kg bodyweight.	Mixture should be given at 6.25mL per 50kg bodyweight.

Horses

COOPERS PANACUR 100 can be administered by a dosing syringe or drenching gun. COOPERS PANACUR 100 has a safety margin in excess of 50 times the recommended dose. COOPERS PANACUR 100 may be safely administered to pregnant mares and working stallions.

Dose Rate Horses: 1mL COOPERS PANACUR 100 per 10kg bodyweight (10mg fenbendazole/kg bodyweight). To control *Strongyloides*

westeri in foals a dose of 5mL per 10kg (50mg fenbendazole/kg bodyweight) is required.

Dose Rate Guide Horses:

Liveweight kg	Dose Volume (mL)	Number of doses per 5L Pack	Liveweight kg	Dose Volume (mL)	Number of doses per 5L Pack
Up to 50	5.0	1000	251-300	30.0	166
51-100	10.0	500	301-350	35.0	142
101-150	15.0	333	351-400	40.0	125
151-200	20.0	250	401-450	45.0	111
201-250	25.0	200	451-500	50.0	100

For animals over 500kg dose at rate of 1mL per 10kg bodyweight.

When used as directed no specific precautions are required other than those normally associated with the careful handling of stock especially weak or pregnant animals. Internal parasites may develop resistance to drenches (anthelmintics). If drench (anthelmintic) failure with this preparation is suspected, seek veterinary confirmation and notify the manufacturer.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

MEAT (CATTLE): DO NOT USE less than 14 days before slaughter for human consumption.

MEAT (HORSES): DO NOT USE less than 28 days before slaughter for human consumption.

MILK: NIL.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): CATTLE: DO NOT USE less than 14 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Coopers Animal Health on 1 800 226 511 or the APVMA website (www.apvma.gov.au/residues/ESI.shtml).

Note: When COOPERS PANACUR 100 is mixed with Fasinex® or Fascicare®, withholding periods for that product will apply. Please refer to the Fasinex® or Fascicare® label for the appropriate withholding period and Trade Advice statements.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126.

Additional Information is listed in the Material Safety Data Sheet.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

Warranty

Intervet Australia Pty Ltd (IAPL) warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

SPECIALIST ADVICE IN EMERGENCY ONLY
COOPERS ANIMAL HEALTH
(A Division of Intervet Australia Pty Ltd)
1800 226 511
ALL HOURS - AUSTRALIA WIDE

COOPERS ANIMAL HEALTH
A division of Intervet Australia Pty Ltd
91-105 Harpin Street, Bendigo East, Vic 3550

STORE below 30°C (Room Temperature). Do not freeze. Store in tightly closed original container in a well-ventilated area. Do not store for prolonged periods in direct sunlight.

APVMA Approval No. 37088/5L/0109

BATCH

EXPIRY

© Registered Trademarks
© Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.
© 2009



5L



B-LA3001149/03