

CAUTION
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

COOPERS®
REVALOR® 400

Growth promotant for grass fed heifers and steers

ACTIVE CONSTITUENTS: EACH IMPLANT (3 PELLETS) CONTAINS: 60 mg TRENBOLONE ACETATE
12 mg 17 β OESTRADIOL

This product contains a palpable marker.

This product must only be supplied in accordance with Division 2 part 4 of the AgVet Regulations.

(Supply of Hormonal Growth Promotants)

For improved growth promotion in grass fed heifers and steers.

One COOPERS REVALOR 400 implanted in growing steers or heifers will normally produce significant liveweight gains when measured at 400 days (see full claim on side panel and leaflet details)

Contents: **1000 implants** (100 foil packed cassettes x 10 implants)

TO BE USED BY SUBCUTANEOUS EAR IMPLANTATION ONLY

1902130A/00

COOPERS®
EST. 1843



CAUTION
KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

COOPERS®
REVALOR® 400

Growth promotant for grass fed heifers and steers

ACTIVE CONSTITUENTS: EACH IMPLANT (3 PELLETS) CONTAINS: 60 mg TRENBOLONE ACETATE
12 mg 17 β OESTRADIOL

This product contains a palpable marker.

This product must only be supplied in accordance with Division 2 part 4 of the AgVet Regulations.

(Supply of Hormonal Growth Promotants)

For improved growth promotion in grass fed heifers and steers.

One COOPERS REVALOR 400 implanted in growing steers or heifers will normally produce significant liveweight gains when measured at 400 days (see full claim on side panel and leaflet details)

Contents: **1000 implants** (100 foil packed cassettes x 10 implants)

TO BE USED BY SUBCUTANEOUS EAR IMPLANTATION ONLY

COOPERS®
EST. 1843



TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on 1800 226 511 before using this product.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 11 26.

DISPOSAL:

DO NOT dispose of unused implants in any way other than surrender to an approved retailer/wholesaler. Dispose of empty cartridge by wrapping with paper and putting in garbage. Discarded needles should be placed in a designated "sharps" container.

STORAGE:

Store below 25°C (Air conditioning).

APMVA Approval No: 48945/51059

REVALOR is recommended for use in heifers and steers. It can be used during the growing or finishing phase and results in improved weight gain due to deposition of added lean meat.

Growth response may vary widely for nutritional or other reasons. Good quality stock and good quality feed with adequate control of parasites and other diseases is a prerequisite for best results. Improved weight gain can be expected 90-100 days post implantation in good quality healthy stock on good nutrition. The increased growth in this period should be maintained when measured at 400 days. Repeat implantation can be carried out during the growing and finishing phase, however, repeat implantation must not be carried out sooner than 70 days. Other products may be more suitable for lot fed cattle.

Distributed by:

COOPERS ANIMAL HEALTH
A division of Intervet Australia Pty Ltd, 91-105 Harpin Street
BENDIGO EAST VIC 3550

Phone: 1800 226 511

Revalor is a Registered Trademark

Batch No.:

Expiry Date:



FL902130A/00

READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT.

DIRECTIONS FOR USE:

To be used ONLY in accordance with the instructions on this label unless authorized under appropriate legislation.

Restraints

FOR USE ONLY in heifers and steers.

DO NOT USE in heifers which are producing or may in the future produce milk or milk products for human consumption.

DO NOT USE this product in any other species of animal.

USE ONLY one implant per heifer or steer.

Implant ONLY beneath the skin in the middle third of the ear.

DO NOT implant this product in any other site.

DO NOT USE this product on HGP free accredited cattle.

It is a breach of AQIS accreditation conditions to use this product on HGP free accredited cattle or to possess this product on any HGP free property (except for feedlots with flexible accreditation). Cattle implanted with this product must be individually identified with an approved ear punch mark.

DO NOT implant sooner than 70 days after last implant.

PRECAUTIONS:

Implanting cattle under 6 weeks of age IS NOT recommended.

Implanting cattle intended for breeding IS NOT recommended as they may experience decreased fertility.

Growth responses may vary widely for nutritional or other reasons. Improved performance may be noted only when conditions are favourable for good growth. Vaginal, rectal and preputial prolapse, increased bulling activity, high tailheads, sunken loins, ventral oedema and udder development may occur as side effects of treatment. To help avoid abscess formation at the site of the implant, the surface of the ear should be thoroughly cleaned and disinfected before implantation.

ADMINISTRATION:

Approved implantation technique is fully described in the enclosed leaflet.

WITHHOLDING PERIOD:

MEAT: 0 days.

MILK: DO NOT USE in heifers which are producing or may in the future produce milk or milk products for human consumption.

