

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



COOPERS®

EASY-DOSE

POUR-ON CATTLE LICE AND FLY TREATMENT

ACTIVE CONSTITUENT: 15.0 g/L DELTAMETHRIN

For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*) and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle. For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses.

Provides protection from *Culicoides brevitarsis* for at least 14 days with effective control of blood feeding *C. brevitarsis* for 6 days post treatment.

CATTLE: ZERO (0) DAYS WITHHOLDING PERIODS FOR MILK & MEAT.

ESI (Export Slaughter Interval): 21 Days.

193479 R3

20 L

COOPERS®
EST. 1843



COOPERS®
EASY-DOSE
POUR-ON CATTLE LICE AND FLY TREATMENT

DIRECTIONS FOR USE

Restraints: NOT TO BE used on horses that may be slaughtered for human consumption.

Contraindications: DO NOT USE on animals severely stressed or in ill health. DO NOT USE on horse's head forward of the ears. DO NOT APPLY as a single spot.

Precautions: Facial skin contact may cause temporary tingling / burning.

DOSE RATE FOR CATTLE: 5 mL per 100 kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Cattle Lice (<i>Bovicola bovis</i>) (<i>Linognathus vituli</i>) (<i>Haematopinus eurysternus</i>)	5 mL	Up to 100 kg	Apply to animals in a race or drainage pen as a single stripe along the backline. Retreat later in the season if reinfestation occurs.
	10 mL	101-200 kg	
	15 mL	201-300 kg	
	20 mL	301-400 kg	
Buffalo Fly (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>) House Fly (<i>Musca domestica</i>) Biting Midge (<i>Culicoides</i> spp.)	25 mL	401-500 kg	Apply to the whole herd – retreatment may be necessary.
	30 mL	501-600 kg	
	Cattle in excess of 600 kg to receive 5 mL per 100 kg liveweight.		

DOSE RATE FOR HORSES: 5 mL per 100 kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Body Weight	Critical Comments
Buffalo Fly (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>)	5 mL	Up to 100 kg	When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice. Apply to the whole herd – retreatment may be necessary.
	10 mL	101-200 kg	
	15 mL	201-300 kg	
	20 mL	301-400 kg	
	25 mL	401-500 kg	
	30 mL	501-600 kg	
	For horses over 600 kg apply 5 mL per 100 kg		

Using the correct measured dose from the squeeze bottle, apply to horses as two single stripes. Apply approximately $\frac{2}{3}$ of measured dose to the midline mane area (from poll to wither) and remaining $\frac{1}{3}$ to midline over the rump.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS (CATTLE): MEAT: Zero (0) days; MILK: Zero (0) days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI) CATTLE: DO NOT USE less than 21 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Coopers Animal Health on 1800 226 511 or the APVMA website (www.apvma.gov.au/residues).

WITHHOLDING PERIODS (HORSES): MEAT: NOT TO BE USED on horses that may be slaughtered for human consumption.

General directions

Cattle: To avoid operator insecticide contact when concurrent treatments are carried out, COOPERS EASY-DOSE should be applied last. Operators must wear gloves.

In a proportion of treated animals, some scurfiness may develop at the site of application. In most cases this scurfiness will disappear within three weeks of application of this product.

Apply with recommended applicator.

The efficacy of this product was not affected by moderate rainfall when tested against cattle lice.

Horses: Trials in horses have shown that treatment with COOPERS EASY-DOSE provides alleviation of fly annoyance related behaviour including: skin flicking, head and tail twitching and feet stamping for up to 7 days. No adverse reactions were reported in horses treated at 5 times the recommended dose rate.

It is recommended the product be allowed to dry thoroughly after application before handling the animal. However, if handling or grooming the animal immediately after application is necessary wear personal protective equipment as indicated in the safety directions on this label.

ENVIRONMENTAL PROTECTION

DANGEROUS TO FISH. Do not contaminate dams, rivers or streams with the pesticide or used containers.

AVOID CONTACT WITH FOOD.

SAFETY DIRECTIONS: Product is poisonous if swallowed. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When using the pour-on wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126. **Additional information is listed in the safety data sheet.**

DISPOSAL: This container can be recycled if it is clean, dry, free of visible residues and has the **drum/MUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drum/MUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

STORAGE: Store below 30°C (room temperature) in original container, tightly closed in a safe place.

COOPERS ANIMAL HEALTH

Intervet Australia Pty Limited
 (trading as MSD Animal Health)
 91-105 Harpin Street,
 Bendigo East VIC 3550
 Phone: 1800 226 511

Warranty

Intervet Australia Pty Limited (IAPL), trading as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

APVMA Approval No.: 54096/52834



GHS Information

Toxic to aquatic life with long lasting effects. Do not eat, drink or smoke when using this product. Avoid release to the environment. **IF SWALLOWED:** Rinse mouth. Do NOT induce vomiting. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and continue rinsing. If skin irritation occurs: Get medical advice. If eye irritation persists: Get medical advice. Not combustible. Use extinguishing media suited to burning materials. Store in a well-ventilated place. Protect from sunlight. Store at temperatures not exceeding 30°C. Store in a dry place. Store in a closed container. For additional information refer to SDS (www.coopersanimalhealth.com.au).

SEE BELOW FOR BATCH AND EXPIRY

® Registered Trademark

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.

20 L



182112 R3

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



COOPERS®

EASY-DOSE

POUR-ON CATTLE LICE AND FLY TREATMENT

ACTIVE CONSTITUENT: 15.0 g/L DELTAMETHRIN

For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*) and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle. For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses.

Provides protection from *Culicoides bevitarsis* for at least 14 days with effective control of blood feeding *C. bevitarsis* for 6 days post treatment.

CATTLE: ZERO (0) DAYS WITHHOLDING PERIODS FOR MILK & MEAT.

ESI (Export Slaughter Interval): 21 Days.

192344 R4

5 L

COOPERS®
EST. 1843

