



Read the enclosed leaflet for full instructions DIRECTIONS FOR USE encing drenching read the label Before com carefully.

Restraints DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption. DO NOT USE in lambs weighing 15 kg or

less. DO NOT USE in calves weighing 100 kg or less.

Contraindications Contraindications This product is contraindicated for use in lambs under six (6) weeks of age or weighing 15 kg or less. This product is contraindicated for use in calves less than 16 weeks of age or weighing 100 kg or less.

Dosage and administration SHAKE WELL BEFORE USE. Dose rates: 1 mL per 10 kg bodyweight by a single oral administration.

SAFETY DIRECTIONS Harmful if inhaled. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent dothing) and elbow-length chemical resistant gloves. Wash hands alter use. Alter each day's use, wash gloves and contaminated dothins. ninated clothing

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT (CATTLE, SHEEP): DO NOT USE less than 21 days before slaughter for human

MILK (CATTLE): DO NOT USE in lactating cows or within 28 days of calving where milk or milk products will be used for human consumption. If cows calve earlier than 28 days after treatment, milk may contain residues. This milk MUST NOT be used for human consumption, or supplied for processing for at least 28 days following treatment. Calves fed this milk should not be slaughtered for human consumption within 21 days. MILK (SHEEP): DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumpt TRADE ADVICE **EXPORT SLAUGHTER INTERVAL (ESI)** CATTLE: DO NOT USE less than 21 days

before slaughter for export. SHEEP: DO NOT USE less than 28 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 1800 226 511 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS

Harmful if inhaled. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbowlength chemical resistant gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. Additional information is listed in the safety data sheet.

For further information contact COOPERS Customer Service on 1800 226 511.

ENVIRONMENTAL PROTECTION

Very toxic to aquatic life. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used containers.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the *drum*MUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State leaislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drum**MUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.



STORAGE

Store below 30°C (room temperature) in the closed original container in a well-ventilated area. Protect from light

Warranty

Intervet Australia Pty Limited (IAPL), trading as MSD Animal Health, warrants that this product is of merchantable auality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

COOPERS ANIMAL HEALTH

Intervet Australia Pty Limited (trading as MSD Animal Health) 91-105 Harpin Street, Bendigo East VIC 3550 Phone: 1800 226 511

APVMA Approval No.: 67327/61112

® Registered trademark ® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered trademark. PANTONE is a registered trademark of Pantone, Inc. © Copyright 2021

COOPERS TRIFECTA is a combination of three active ingredients including abamectin (a member of the Macrocyclic Lactone (ML) drench family); oxfendazole (a member of the Benzimidazole (white) drench family) and levamisole hydrochloride (an Imidazothiazole and member of the clear drench family). It also contains two minerals including selenium (as sodium selengte) and cobalt (as cohalt FDTA)

COOPERS TRIFECTA provides treatment and control of the following roundworms, sensitive to at least one of the three active ingredients in this product.

SHEED

Adult gastro-intestinal roundworms

- Barber's Pole Worm (Haemonchus contortus) Small Brown Stomach Worm (Teladorsaaia circumcincta)
- Stomach Hair Worm (Trichostrongylus axei)
- Black Scour Worm (Trichostrongylus colubriformis[†]; T vitrinus) (⁺ including *T. colubriformis* resistant to a combination of
- benzimidazoles and levamisole) Small Intestinal Worm (Cooperia curticei; C. oncophora)
- Thin Necked Intesting Worm (Nematodirus filicollis: N. spathiger; N. abnormalis)
- Hookworm (Bunostomum trigonocephalum)
- Intestinal Threadworm (*Strongyloides papillosus*)
- Large Bowel Worm (*Oesophagostomum venulosum*)
- Nodule worm (*Oesophagostomum columbianum*) • Large Mouthed Bowel Worm (Chabertia ovina)
- Whipworm (Trichuris ovis)

CATTIF

Adult gastro-intestinal roundworms

Barber's Pole Worm (Haemonchus placei)

- Small Brown Stomach Worm (Ostertagia ostertagi)
- Stomach Hair Worm (Trichostrongylus axei)

PRODUCT LABEL | Trifecta 2.5L Leaflet

- Black Scour Worm (Trichostrongylus longispicularis) • Small Intestinal Worm (Cooperia oncophera; C. pectinata; C. punctata)
- Thin Necked Intestinal Worm (Nemtaodirus helvetianus)
- Hookworm (Bunostomum phlebotomum)
- Capillarids (*Capillaria bovis*)
- Nodule Worm (*Oesonhaaostomum radiatum*)

Resistance may develop to any drench. Ask your local veterinarian or animal health advisor for recommended parasite management practices for your area to reduce resistance development. It is advisable that a drench resistance test should be conducted before any drench is used

DIRECTIONS FOR USE

Before commencing drenching read the label carefully.

Restraints

DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption. DO NOT USE in lambs weighing 15 kg or less. DO NOT USE in calves weighing 100 kg or less.

Contraindications

This product is contraindicated for use in lambs under six (6) weeks of age or weighing 15 kg or less.

This product is contraindicated for use in calves less than 16 weeks of age or weigh 100 ka or less.

This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by

other means (such as pasture top dressing, vaccine,

pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.

Precautions

Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress. Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching. Recommended dose should not be exceeded. Selenium can have toxic effects, particularly in cattle and sheep with liver damage including cattle and sheep that araze in areas where plants containing pyrrolizidine alkaloids are prevalent. Safety in breeding cattle and sheep of either sex and in pregnant or lactating cows or ewes has not been established

Dosage and administration SHAKE WELL BEFORE USE.

COOPERS TRIFECTA should be administered orally using accurate drenching equipment. Clean equipment thoroughly before and after use. Always dose on a liveweight basis, do not underdose. The heaviest animal in a group should be weighed.

Dose the group to the heaviest animal by liveweight (as ewes, wethers, rams, lambs, cows, bulls, calves), do not underdose. For cattle, where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Check the accuracy of drenching equipment before and during drenching procedures.

Re-Treatment Interval (SHEEP/CATTLE): DO NOT retreat animals for 28 days after last treatment.

Dose rates: 1 mL per 10 kg bodyweight by a single oral administration



7 I - 80 kg Over 80 kg	8 mL 1 mL/10 kg	312	
CATTLE LIVEWEIGHT 100 kg or less	DOSE Do not treat	2.5 L PACK TREATS	
101 - 150 kg	15 mL	166	
151 - 200 kg	20 mL	125	
201 - 250 kg	25 mL	100	
251 - 300 kg	30 mL	83	
301 - 350 kg	35 mL	71	
351 - 400 kg	40 mL	62	
401 - 450 kg	45 mL	55	
451 - 500 kg	50 mL	50	
501 - 550 kg	55 mL	45	
551 - 600 kg	60 mL	41	

65 mL

1 mL/10 kg

38

601 - 650 kg

Over 650 kg

DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption. DO NOT USE in lambs weighing 15 kg or less. DO NOT USE in calves weighing 100 kg or less.

This product is contraindicated for use in lambs under six (6) weeks of age or weighing 15 kg or less. This product is contraindicated for use in calves less than 16 weeks of age or weighing 100 kg or less.

Harmful if inhaled. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands alter use. Alter each day's use, wash gloves and contaminated clothing. FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. Additional information is listed in the safety data sheet.

Read the enclosed leaflet for full instructions.

DIRECTIONS FOR USE

Before commencing drenching read the label carefully.

RESTRAINTS

Contraindications

Dosage and administration

SHAKE WELL BEFORE USE. **Dose rates:** 1 mL per 10 kg bodyweight by a single oral administration.

SAFETY DIRECTIONS