

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

FLEXOLT®

Oral Lice Treatment for sheep with Any Length of Wool

ACTIVE CONSTITUENT: 10 g/L FLURALANER

For the control of isoxazoline-susceptible lice (*Bovicola ovis*), including strains resistant to synthetic pyrethroids and insect growth regulators, on sheep and lambs with any length of wool.

20 L



167398 R1

FLEXOLT®

Oral Lice Treatment for sheep with Any Length of Wool

Fluralaner is a molecule from the isoxazoline family of chemicals and is a novel compound that has shown to be highly efficacious against sheep body lice (*Bovicola ovis*). It works systemically to kill lice in sheep and lambs no matter the length of wool.

FLEXOLT has been shown to be effective on a wide range of sheep across a large number of studies including off-shears sheep, shorn sheep up to 7 days off-shears, unshorn lambs, sheep with short and long wool, and off-shears and long wool ewes with lambs at foot.

FLEXOLT has been shown to be safe at 5x the maximum recommended dose rate (15 mg fluralaner per kg body weight) and safe for the treatment of young lambs as small as 6 kg body weight.

Lice control is not affected by rainfall before or after treatment.

Resistance may develop to any chemical.

DIRECTIONS FOR USE

Restraints

DO NOT USE in lambs less than 6 kg body weight. DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

Re-treatment interval: DO NOT re-treat animals for 54 days after last treatment.

Precautions

A proportion of sheep may cough following treatment. This passing response is of no consequence. Safety in ewes during mating has not been established.

Dosage and administration

This product is for oral application only, do not give topically or parenterally.

FLEXOLT is safe for use in breeding rams and pregnant and lactating ewes, however, safety in ewes during mating has not been established.

Check the accuracy of the applicator before and during drenching procedure.

The recommended dose rate is 0.15 mL per kg body weight (equivalent to 1.5 mg fluralaner per kg body weight).

Dose rate: 1.5 mL per 10 kg body weight.

The following table may be used as a guide:

Body weight (kg)	Dose (mL)
6 - 10	1.5
11 - 16	2.5
17 - 20	3.0
21 - 30	4.5
31 - 40	6.0
41 - 50	7.5
51 - 60	9
61 - 70	11
71 - 80	12
81 - 90	14
91 - 100*	15

* Add 3 mL for each 20 kg above 100 kg.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by body weight in each group (ewes, wethers, rams, lambs). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

It is essential that every sheep is treated to achieve lice control – it is recommended that as sheep are treated, they are marked so any untreated sheep can be identified and treated. Do not leave any sheep in the flock untreated.

Critical use comments

- Administer orally using accurate drench equipment or syringes.
- DO NOT mix treated sheep with untreated sheep.
- Avoid treating ewes less than 4 weeks before lambing commences or while ewes are lambing, as live lice can persist and infect the lambs. Ewes and lambs can be treated together after lambing has finished.

General directions

FLEXOLT is a short acting, non-residual product. Maximum lice kill at treatment and strict quarantine conditions for treated sheep are necessary for effective lice control. It is recommended that a thorough muster be conducted so that no untreated sheep remain. Boundary fences should be checked to ensure animals are contained and stray sheep are excluded. Treated sheep must be kept separate from lice infested sheep or sheep of unknown lice status. For effective lice control, every sheep on the property should be treated at the same time.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.

WOOL HARVESTING INTERVAL: Zero (0) days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 54 days before slaughter for export. Before using this product, confirm the current ESI from Coopers Animal Health on 1800 226 511 or the APVMA website (apvma.gov.au/residues).

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the product for use and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.

ADDITIONAL USER SAFETY

Additional information is listed in the safety data sheet (SDS).

ENVIRONMENTAL PROTECTION

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the **drumMUSTER** logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container.



If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 30°C (room temperature) in the tightly closed, original container in a cool dry place. Do not store for prolonged periods in direct sunlight.

Warranty

Intervet Australia Pty Limited (IAPL), trading as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

COOPERS ANIMAL HEALTH

Intervet Australia Pty Limited
(trading as MSD Animal Health)
91-105 Harpin Street,
Bendigo East VIC 3550
Phone: 1800 226 511

APVMA Approval No.: 91565/132669

SEE BELOW FOR BATCH AND EXPIRY

© Registered trademark.

© Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a registered trademark. PANTONE is a registered trademark of Pantone, Inc. Copyright © 2023 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.

GHS Information

Wear eye / face protection.

For additional information refer to SDS (www.coopersanimalhealth.com.au).



Batch
Expiry

Imprint area

20 L



Scan the QR code for more information.

167650 R2