

CAUTION


KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

COOPERS®

Exzolt®

POUR-ON FOR CATTLE

ACTIVE CONSTITUENT: 50 g/L FLURALANER
ALSO CONTAINS: 350 g/L 2-PYRROLIDONE (SOLVENT)



Coopers Exzolt Pour-on for Cattle is for the control of isoxazoline-susceptible strains of Cattle Tick (*Rhipicephalus (Boophilus) microplus*) including dieldrin, organophosphate, synthetic pyrethroid, amidine, macrocyclic lactone and fluzuron resistant ticks. Controls Bush Tick (*Haemaphysalis longicornis*) and Buffalo Fly (*Haematobia irritans exigua*) including synthetic pyrethroid resistant strains.

Coopers Exzolt Pour-on for Cattle controls:

- Cattle Tick for 56 days
- Bush Tick for 21 days
- Buffalo Fly for 21 days

Coopers Exzolt Pour-on for Cattle is a next generation ectoparasiticide. It contains Fluralaner, a parasiticide belonging to the isoxazoline family of chemicals, which works systemically as well as by contact to control ectoparasites.

Resistance may develop to any chemical.

DIRECTIONS FOR USE

Restraints

DO NOT USE in cows which are producing or may in the future produce milk that may be used or processed for human consumption except replacement dairy heifers prior to first mating.

Re-treatment interval: DO NOT re-treat animals for 88 days after last treatment.

Precautions

Reproductive safety in breeding bulls has not been evaluated. Safety has not been demonstrated in calves younger than 6 months.

Side effects

Topical treatment with **Coopers Exzolt Pour-on for Cattle** is generally well tolerated and no serious adverse reactions have occurred in the studies. Mild and transient local skin reactions may occur, which do not affect the general health of the animals and do not require specific treatment. If a skin reaction does occur, it is cosmetic in nature and will typically resolve within 41 days after treatment.

Dosage and administration

Administer topically in a narrow strip along the midline of the back from the withers to the base of the tail. Cattle should be dry at treatment. Use in well-ventilated areas or outdoors.


Coopers Exzolt Pour-on for Cattle is safe for use in breeding and pregnant cows, however, reproductive safety in breeding bulls has not been evaluated. Keep treated animals separated from untreated animals.

The recommended dose rate is 1 mL/20 kg body weight (2.5 mg Fluralaner per kg body weight).

Dose rate: 1 mL per 20 kg body weight.

The following table may be used as a guide:

Body weight (kg)	Dose volume (mL)	5 L Pack treats (Cattle)
201 – 250	12.5	400
251 – 300	15.0	333
301 – 350	17.5	285
351 – 400	20.0	250
401 – 450	22.5	222
451 – 500	25.0	200
501 – 550	27.5	181
551 – 600	30.0	166
601 – 650*	32.5	153



* Add 2.5 mL for each 50 kg above 650 kg.

A representative sample of animals should be weighed before treatment either with scales or a weighband.

Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not under dose.

Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing.

Apply only with the recommended applicator and draw off tube. Clean applicator with warm soapy water then flush with clean water and allow to dry before next use.

General directions

The efficacy of **Coopers Exzolt Pour-on for Cattle** is not adversely affected by heavy rainfall after treatment. Study data demonstrated that artificial rain applied by inverted sprinklers (equivalent to rainfall of 22-23 mm in a storm lasting approximately 20 minutes) 6, 12, or 24 hours following treatment did not adversely affect efficacy.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 38 days before slaughter for human consumption.

MILK: DO NOT USE in cows which are producing or may in the future produce milk that may be used or processed for human consumption except replacement dairy heifers prior to first mating.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 88 days before slaughter for export. Before using this product, confirm the current ESI from MSD Animal Health on 1800 226 511 or the APVMA website (apvma.gov.au/residues).

SAFETY DIRECTIONS

Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

Re-handling interval: DO NOT handle treated animals until product has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. If swallowed, do NOT induce vomiting. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes wash out immediately with water.

ADDITIONAL USER SAFETY

Additional information is listed in the safety data sheet (SDS).

ENVIRONMENTAL PROTECTION

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers. Toxic to dung fauna. However, use of the product as directed is not expected to have long-term adverse effects on dung fauna or dung degradation.

DISPOSAL

Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the **drumMUSTER** logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.



STORAGE

Store below 30°C (room temperature).

Warranty

Intervet Australia Pty Limited (IAPL), trading as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

Intervet Australia Pty Limited
(trading as MSD Animal Health)
91-105 Harpin Street,
Bendigo East VIC 3550
Phone: 1800 226 511

APVMA Approval No.: 92557/135894.

® Registered trademark.

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a registered trademark. PANTONE is a registered trademark of Pantone, Inc. Copyright © 2024 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.

GHS Information

Flammable liquid and vapour. May cause drowsiness or dizziness.

May damage fertility. May damage the unborn child. Keep away

from heat, hot surfaces, sparks, open flames and other ignition

sources. No smoking. Avoid breathing mist or vapours. If on skin

or hair, take off immediately all contaminated clothing. If inhaled,

remove person to fresh air and keep comfortable for breathing. If in

eyes, rinse cautiously with water for several minutes. Remove contact

lenses, if present and easy to do. Continue rinsing. If eye irritation

persists, get medical advice.



Scan the QR code for
more information.



MSD
Animal Health