

POISON


KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

COOPERS®

EASY-DOSE**POUR-ON CATTLE LICE
AND FLY TREATMENT****ACTIVE CONSTITUENT: 15.0 g/L DELTAMETHRIN**

For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*), and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle.

For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses. Provides protection from *Culicoides brevitarsus* for at least 14 days with effective control of blood feeding *C.brevitarsus* for 6 days post treatment.

GUN PACK**PICTURE OF 5L PACK + GUN]****5 Litres****COOPERS LOGO**

File Name: 54096_52834-RLP_V2		Version: 2	Page 1 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

[CATTLE, HORSE PICTURE]

COOPERS®

EASY-DOSE

POUR-ON CATTLE LICE AND FLY TREATMENT

GUN PACK

DIRECTIONS FOR USE:

For details of Directions for Use refer to container label.

DOSE RATE FOR CATTLE: 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Cattle Lice (<i>Bovicola bovis</i>) (<i>Linognathus vituli</i>) (<i>Haematopinus eurysternus</i>)	5mL 10mL 15mL 20mL 25mL 30mL	Up to 100kg 101-200kg 201-300kg 301-400kg 401-500kg 501-600kg	Apply to animals in a race or drainage pen as a single stripe along the backline. Retreat later in the season if reinfestation occurs.
Buffalo Fly (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>) House Fly (<i>Musca domestica</i>) Biting Midge (<i>Culicoides</i> sp.)		Cattle in excess of 600kg To receive 5mL per 100kg liveweight.	Apply to the whole herd – re-treatment may be necessary.

DOSE RATE FOR HORSES: 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Body Weight	Critical Comments
Buffalo Fly \ (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>)	5mL 10mL 15mL 20mL 25mL 30mL	Up to 100kg 101-200kg 201-300kg 301-400kg 401-500kg 501-600kg For horses over 600kg apply 5mL per 100kg	When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice. Apply to the whole herd – retreatment may be necessary.

Using the correct measured dose from the squeeze bottle, apply to horses as two single stripes. Apply approximately 2/3 of measured dose to the midline mane area (from poll to wither) and remaining 1/3 to midline over the rump.

SPECIALIST ADVICE IN EMERGENCY ONLY

COOPERS ANIMAL HEALTH
(A Division of Intervet Australia Pty Ltd)
1 800 033 461
ALL HOURS – AUSTRALIA WIDE

SAFETY DIRECTIONS

Product is poisonous if swallowed. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When using the pour-on wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Additional information is listed in the Material Safety Data Sheet.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

COOPERS ANIMAL HEALTH

A division of Intervet Australia Pty Ltd
91-105 Harpin Street, Bendigo East VIC 3550

STORE below 30°C (Room Temperature) in original container, tightly closed in a safe place.
APVMA Approval No. 54096/52834

BATCH**EXPIRY**

® Registered Trademark

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.

© 2009

File Name: 54096_52834-RLP_V2		Version: 2	Page 3 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

[PICTURE]

COOPERS®
EASY-DOSE

**POUR-ON CATTLE LICE
AND FLY TREATMENT**

GUN PACK

**5 Litres
EASY-DOSE**

[PICTURE OF 5L CONTAINER]

**EASY-DOSE
APPLICATOR**

[PICTURE OF APPLICATOR]

[PICTURE]

COOPERS®
EASY-DOSE
POUR-ON CATTLE LICE
AND FLY TREATMENT
GUN PACK

COOPERS®
EASY-DOSE
POUR-ON CATTLE LICE
AND FLY TREATMENT
GUN PACK

[PICTURE]

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

[Cattle Picture]

Coopers®

EASY-DOSE
POUR-ON CATTLE LICE AND FLY TREATMENT

ACTIVE CONSTITUENT: 15.0 g/L DELTAMETHRIN

For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*), and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle.

For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses.

Provides protection from *Culicoides brevitarsus* for at least 14 days with effective control of blood feeding *C.brevitarsus* for 6 days post treatment.

20 Litres
25 Litres

COOPERS LOGO

CATTLE: ZERO (0) DAYS WITHHOLDING PERIOD FOR MILK & MEAT.
ESI (Export Slaughter Interval): 21 Days.

File Name: 54096_52834-RLP_V2		Version: 2	Page 6 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

POISON**KEEP OUT OF REACH OF CHILDREN****READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY****[Cattle Picture]**

Coopers®

EASY-DOSE**POUR-ON CATTLE LICE
AND FLY TREATMENT****ACTIVE CONSTITUENT: 15.0 g/L DELTAMETHRIN**

For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*), and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle.

For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses.

Provides protection from *Culicoides brevitarsus* for at least 14 days with effective control of blood feeding *C.brevitarsus* for 6 days post treatment.

**2 Litres
5 Litres
6 Litres****COOPERS LOGO****CATTLE: ZERO (0) DAYS WITHHOLDING PERIOD FOR MILK & MEAT
ESI (Export Slaughter Interval): 21 Days.**

File Name: 54096_52834-RLP_V2		Version: 2	Page 7 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

Coopers®

EASY-DOSE**POUR-ON CATTLE LICE AND FLY TREATMENT****DIRECTIONS FOR USE:****RESTRAINTS****NOT TO BE USED on horses that may be slaughtered for human consumption.****CONTRAINDICATIONS:** DO NOT USE on animals severely stressed or in ill health. DO NOT USE on horse's head forward of the ears. DO NOT APPLY as a single spot.**PRECAUTION:** Facial skin contact may cause temporary tingling / burning.**DOSE RATE FOR CATTLE:** 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Cattle Lice (<i>Bovicola bovis</i>) (<i>Linognathus vituli</i>) (<i>Haematopinus eurysternus</i>)	5mL 10mL 15mL 20mL 25mL 30mL	Up to 100kg 101-200kg 201-300kg 301-400kg 401-500kg 501-600kg	Apply to animals in a race or drainage pen as a single stripe along the backline. Retreat later in the season if reinfestation occurs.
Buffalo Fly (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>) House Fly (<i>Musca domestica</i>) Biting Midge (<i>Culicoides</i> sp.)	Cattle in excess of 600kg to receive 5 mL per 100 kg liveweight.		Apply to the whole herd - re-treatment may be necessary.

DOSE RATE FOR HORSES: 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Body Weight	Critical Comments
Buffalo Fly \ (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>)	5mL 10mL 15mL 20mL 25mL 30mL For horses over 600kg apply 5mL per 100kg	Up to 100kg 101-200kg 201-300kg 301-400kg 401-500kg 501-600kg	When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice. Apply to the whole herd – retreatment may be necessary.

Using the correct measured dose from the squeeze bottle, apply to horses as two single stripes. Apply approximately 2/3 of measured dose to the midline mane area (from poll to wither) and remaining 1/3 to midline over the rump.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

File Name: 54096_52834-RLP_V2		Version: 2	Page 8 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

WITHHOLDING PERIODS: CATTLE: MEAT ZERO (0) DAYS; MILK ZERO (0) DAYS.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Coopers Animal Health on 1800 226 511 or the APVMA website (www.apvma.gov.au/residues/ESI.html).

MEAT WITHHOLDING PERIOD (HORSES): NOT TO BE USED on horses intended for human consumption.

GENERAL INSTRUCTIONS

Cattle:

To avoid operator insecticide contact when concurrent treatments are carried out, COOPERS EASY-DOSE should be applied last. Operators must wear gloves.

In a proportion of treated animals, some scurfiness may develop at the site of application. In most cases this scurfiness will disappear within three weeks of application of this product.

Apply with recommended applicator.

The efficacy of this product was not affected by moderate rainfall when tested against cattle lice.

Horses: Trials in horses have shown that treatment with COOPERS EASY-DOSE provides alleviation of fly annoyance related behaviour including: skin flicking, head and tail twitching and feet stamping for up to 7 days. No adverse reactions were reported in horses treated at 5 times the recommended dose rate.

It is recommended the product be allowed to dry thoroughly after application before handling the animal. However, if handling or grooming the animal immediately after application is necessary wear personal protective equipment as indicated in the safety directions on this label.

For further information contact Coopers Customer Service on 1 800 226511

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

DANGEROUS TO FISH. Do not contaminate dams, rivers or streams with the pesticide or used containers.

AVOID CONTACT WITH FOOD.

SAFETY DIRECTIONS

Product is poisonous if swallowed. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When using the pour-on wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

Additional information is listed in the Material Safety Data Sheet.

File Name: 54096_52834-RLP_V2		Version: 2	Page 9 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

Warranty

Intervet Australia Pty Ltd (IAPL) warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

SPECIALIST ADVICE IN EMERGENCY ONLY

COOPERS ANIMAL HEALTH
(A Division of Intervet Australia Pty Ltd)
1 800 226511
ALL HOURS – AUSTRALIA WIDE

**DRUMMUSTER
ELIGIBLE
CONTAINER**

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

COOPERS ANIMAL HEALTH

A division of Intervet Australia Pty Ltd
91-105 Harpin Street, Bendigo East VIC 3550

STORE below 30°C (Room Temperature) in original container, tightly closed in a safe place.

APVMA Approval No. 54096/52834

FOR BATCH AND EXPIRY SEE BELOW

® Registered Trademark

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.

© 2009

2L/5L/6L/20L/25L

[Batch and Expiry]

File Name: 54096_52834-RLP_V2		Version: 2	Page 10 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

[CATTLE PICTURE]

Coopers®

EASY-DOSE
POUR-ON CATTLE LICE
AND FLY TREATMENT

ACTIVE CONSTITUENT: 15.0 g/L DELTAMETHRIN

For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*), and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle.

For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses.

Provides protection from *Culicoides brevitarsus* for at least 14 days with effective control of blood feeding *C.brevitarsus* for 6 days post treatment.

1 Litre

COOPERS LOGO

CATTLE: ZERO (0) DAYS WITHHOLDING PERIOD FOR MILK & MEAT
ESI (Export Slaughter Interval): 21 Days.

File Name: 54096_52834-RLP_V2		Version: 2	Page 11 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

COOPERS®**EASY-DOSE****POUR-ON CATTLE LICE AND FLY TREATMENT**

SAFETY DIRECTIONS: Product is poisonous if swallowed. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When using the pour-on wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.

FIRST AID:

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126. Additional information is listed in the material safety data sheet.

WARRANTY

Intervet Australia Pty Ltd (IAPL) warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

For further information contact Coopers Customer Service on 1-800-226-511.

SPECIALIST ADVICE IN EMERGENCY ONLY

COOPERS ANIMAL HEALTH
(A Division of Intervet Australia Pty Ltd)

1 800 226511

ALL HOURS - AUSTRALIA-WIDE

Dispose of empty container by wrapping with paper and putting in garbage.

COOPERS ANIMAL HEALTH

A division of Intervet Australia Pty Ltd
91-105 Harpin Street, Bendigo East VIC 3550

Store below 30°C (Room Temperature) in original container, tightly closed in a safe place.

For Batch and Expiry See Below

APVMA Approval No. 54096/52834

® Registered Trademark

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.

© 2009

1L

[Batch and Expiry]

File Name: 54096_52834-RLP_V2		Version: 2	Page 12 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

TO OPEN PEEL HERE ⇨

COOPERS®

EASY-DOSE**Pour-on Cattle Lice and Fly Treatment**

For control of Stable Fly (*Stomoxys calcitrans*), House Fly (*Musca domestica*) and Lice (*Bovicola bovis*, *Linognathus vituli* and *Haematopinus eurysternus*) and synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) on beef and dairy cattle.

For control of synthetic pyrethroid susceptible Buffalo Fly (*Haematobia irritans exigua*) for up to 3 weeks and for control of Stable Fly (*Stomoxys calcitrans*) for up to 1 week on horses.

Provides protection from *Culicoides brevitarsus* for at least 14 days with effective control of blood feeding *C. brevitarsus* for 6 days post treatment.

READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT.

SAFETY DIRECTIONS: Product is poisonous if swallowed. Will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When using the pour-on wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.

FIRST AID:

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126. Additional information is listed in the material safety data sheet.

Dispose of empty container by wrapping with paper and putting in garbage.

COOPERS ANIMAL HEALTH

A division of Intervet Australia Pty Ltd
91-105 Harpin Street, Bendigo East VIC 3550

Store below 30°C (Room Temperature) in original container, tightly closed in a safe place.

For Batch and Expiry see below

APVMA Approval No. 54096/52834

® Registered Trademark

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.

© 2009

1L

[Batch and Expiry will be screenprinted directly onto container]

File Name: 54096_52834-RLP_V2		Version: 2	Page 13 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			

DIRECTIONS FOR USE:**RESTRAINTS****NOT TO BE USED** on horses that may be slaughtered for human consumption.**CONTRAINDICATIONS:** DO NOT USE on animals severely stressed or in ill health. DO NOT USE on horse's head forward of the ears. DO NOT APPLY as a single spot.**PRECAUTION:** Facial skin contact may cause temporary tingling / burning.**DOSE RATE FOR CATTLE:** 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Cattle Lice (<i>Bovicola bovis</i>) (<i>Linognathus vituli</i>) (<i>Haematopinus eurysternus</i>)	5mL 10mL 15mL 20mL 25mL 30mL	Up to 100kg 101-200kg 201-300kg 301-400kg 401-500kg 501-600kg	Apply to animals in a race or drainage pen as a single stripe along the backline. Retreat later in the season if reinfestation occurs.
Buffalo Fly (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>) House Fly (<i>Musca domestica</i>) Biting Midge (<i>Culicoides</i> sp.)	Cattle in excess of 600kg to receive 5 mL per 100 kg liveweight.		Apply to the whole herd – re-treatment may be necessary.

DOSE RATE FOR HORSES: 5mL per 100kg liveweight.

The following table may be used as a guide.

Pests Controlled	Rate	Body Weight	Critical Comments
Buffalo Fly \ (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>)	5mL 10mL 15mL 20mL 25mL 30mL For horses over 600kg apply 5mL per 100kg	Up to 100kg 101-200kg 201-300kg 301-400kg 401-500kg 501-600kg	When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice. Apply to the whole herd – retreatment may be necessary.

Using the correct measured dose from the squeeze bottle, apply to horses as two single stripes. Apply approximately 2/3 of measured dose to the midline mane area (from poll to wither) and remaining 1/3 to midline over the rump.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS: CATTLE: MEAT ZERO (0) DAYS; MILK ZERO (0) DAYS.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Coopers Animal Health on 1800 226 511 or the APVMA website (www.apvma.gov.au/residues/ESI.html).

MEAT WITHHOLDING PERIOD (HORSES): NOT TO BE USED on horses intended for human consumption.

GENERAL INSTRUCTIONS

Cattle:

To avoid operator insecticide contact when concurrent treatments are carried out, COOPERS EASY-DOSE should be applied last. Operators must wear gloves.

In a proportion of treated animals, some scurfiness may develop at the site of application. In most cases this scurfiness will disappear within three weeks of application of this product.

Apply with recommended applicator.

The efficacy of this product was not affected by moderate rainfall when tested against cattle lice.

Horses: Trials in horses have shown that treatment with COOPERS EASY-DOSE provides alleviation of fly annoyance related behaviour including: skin flicking, head and tail twitching and feet stamping for up to 7 days. No adverse reactions were reported in horses treated at 5 times the recommended dose rate.

It is recommended the product be allowed to dry thoroughly after application before handling the animal. However, if handling or grooming the animal immediately after application is necessary wear personal protective equipment as indicated in the safety directions on this label.

For further information contact Coopers Customer Service on 1 800 226511.

File Name: 54096_52834-RLP_V2		Version: 2	Page 15 of 15
Prepared By: J.Howlett			
Preparation Date: 28/02/2011			
Last Revised: 15/02/2011			