APVMA Approval No.: 58656/20I/0610 Information Centre. Phone 13 11 26. If poisoning occurs, contact a doctor or Poisons FIRST AID:

1800 226 511 before using this prodeut. advice on the ESI, contact the manufacturer on product does not have an ESI established. For EXPORT SLAUGHTER INTERVAL (ESI): This HADE ADVICE:

milk products for human consumption. producing or may in the future produce milk or MILK: DO NOT use in heifers which are JIN :IA HM

> **МІТННОГДІИ**С РЕВІОДЅ: described in the enclosed leaflet.

Approved implantation technique is fully

:NOITARTSINIMGA

implantation. thoroughly cleaned and disinfected before the implant, the surface of the ear should be To help avoid abscess formation at the site of occur as side effects of treatment. ventral oedema and udder development may bulling activity, high tailheads, sunken loins, Vaginal, rectal and preputial prolapse, increased conditions are tayourable for good growth. bertormance may be noted only when nutritional or other reasons, improved Growth responses may vary widely tor experience decreased fertility. implanting cattle intended for preeding is NOT recommended as they may Implanting cattle under 6 weeks of age is NOT recommended. PRECAUTIONS:

nast implant.

DO NOT implant sooner than 70 days after approved ear punch mark. broduct must be individually identified with an tree property (except for feedlots with flexible accreditation). Cattle implanted with this cattle or to possess this product on any HGP to use this product on HGP free accredited It is a breach of AQIS accreditation conditions accredited cattle.

DO NOT USE this product on HGP free DO NOT implant this product in any other site.

third of the ear. Implant ONLY beneath the skin in the middle USE ONLY one implant per heiter and steer. species of animal.

DO NOT USE this product in any other products for human consumption. may in the future produce milk or milk DO NOT use in heifers which are producing or FOR USE ONLY in non breeding cattle. Restraints

authorised under appropriate legislation. To be used ONLY in accordance with the instructions on this label unless

DIRECTIONS FOR USE:

READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT.

Batch No.: Expiry Date:

DISPOSAL:

DO NOT dispose of unused implants in any way other than surrender to an approved retailer/wholesaler.

Dispose of empty cartridges by wrapping with paper and putting in garbage Discarded needles should be placed in a designated 'sharps' container.

STORAGE: Store below 25°C (Air conditioning).

Distributed by:

COOPERS ANIMAL HEALTH

A Division of Intervet Australia Pty Ltd 91-105 Harpin Street, BENDIGO EAST, VIC 3550

Phone: 1800 226 511

Revalor® is a Registered Trademark

113787 CBW880 03



KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY



Growth promotant for non breeding cattle



ACTIVE CONSTITUENTS: EACH IMPLANT (4 PELLETS) CONTAINS: 80 mg TRENBOLONE ACETATE 16 mg 17 β OESTRADIOL

This product contains a palpable marker.

This product must only be supplied in accordance with Division 2 part 4 of the AgVet Regulations (Supply of Hormonal Growth Promotants)

For improved growth promotion in non breeding cattle.

Contents: 20 (2 cassettes x 10 implants)

TO BE USED BY SUBCUTANEOUS

EAR IMPLANTATION ONLY

Scale: A4 = 100%

Revalor® is recommended for use in non breeding cattle. It can be used during the growing or finishing phase and results in improved weight gain due to deposition of added lean meat. Repeat implantation can be carried out during the growing and finishing phase, however, it is recommended that repeat implantation is not carried out sooner than 70 days and that there is adequate nutrition to ensure a growth response. Growth response may vary widely for nutritional or other reasons. Good quality feed is a prerequisite for best results whether on pasture, supplement or lot feeding. Maximum growth rate requires good quality stock with adequate control of parasites and other diseases.





EAR IMPLANTATION ONLY TO BE USED BY SUBCUTANEOUS Contents: 100 (10 cassettes x 10 implants) For improved growth promotion in non breeding cattle.

(Supply of Hormonal Growth Promotants) This product must only be supplied in accordance with Division 2 part 4 of the AgVet Regulations This product contains a palpable marker.

16 mg 17 ß OESTRADIOL ACTIVE CONSTITUENTS: EACH IMPLANT (4 PELLETS) CONTAINS: 80 mg TRENBOLONE ACETATE

Growth promotant for non breeding cattle

BEAVIOU® FLEX COOFERS®

FOR AUIMAL TREATMENT ONLY KEEP OUT OF REACH OF CHILDREN **NOITUAD**

CAUTION **KEEP OUT OF REACH OF CHILDREN** FOR ANIMAL TREATMENT ONLY

COOPERS® ALOR® FLEX

Growth promotant for non breeding cattle

ACTIVE CONSTITUENTS: EACH IMPLANT (4 PELLETS) CONTAINS: 80 mg TRENBOLONE ACETATE 16 mg 17 ß OESTRADIOL

This product contains a palpable marker.

This product must only be supplied in accordance with Division 2 part 4 of the AgVet Regulations (Supply of Hormonal Growth Promotants)

For improved growth promotion in non breeding cattle.

Contents: 100 (10 cassettes x 10 implants)

TO BE USED BY SUBCUTANEOUS EAR IMPLANTATION ONLY

103111 CBW880 03

READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT.

DIRECTIONS FOR USE: To be used ONLY in accordance with the instructions on this label unless authorised under appropriate legislation

Restraints

FOR USE ONLY in non breeding cattle.

DO NOT use in heifers which are producing or may in the future produce milk or milk products for

DO NOT USE this product in any other species of animal.

USE ONLY one implant per heifer and steer. Implant ONLY beneath the skin in the middle third of

DO NOT implant this product in any other site. DO NOT USE this product on HGP free accredited cattle.

It is a breach of AQIS accreditation conditions to use this product on HGP free accredited cattle or to possess this product on any HGP free property (except for feedlots with flexible accreditation). Cattle implanted with this product must be individually identified with an approved

DO NOT implant sooner than 70 days after last implant.

PRECAUTIONS: Implanting cattle under 6 weeks of age is NOT recommended. Implanting cattle intended for breeding is NOT recommended as they may experience decreased fertility. Growth responses may vary widely for nutritional or other reasons. Improved performance may be noted only when conditions are favourable for good growth. Vaginal, rectal and preputial prolapse, increased bulling activity, high tailheads, sunken loins, ventral oederna and udder development may occur as side effects of treatment.

To help avoid abscess formation at the site of the implant, the surface of the ear should be thoroughly cleaned and disinfected before implantation.

COOPERS ANIMAL HEALTH

A Division of Intervet Australian Pty Ltd 91-105 Harpin Street, BENDIGO EAST, VIC 3550 Phone: 1800 226 211

revalor[®] is a Registered Trademark





CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY

COOPERS®

ALOR® FLEX

Growth promotant for non breeding cattle

ACTIVE CONSTITUENTS: EACH IMPLANT (4 PELLETS) CONTAINS: 80 mg TRENBOLONE ACETATE 16 mg 17 β OESTRADIOL

This product contains a palpable marker.

This product must only be supplied in accordance with Division 2 part 4 of the AgVet Regulations (Supply of Hormonal Growth Promotants)

For improved growth promotion in non breeding cattle.

Contents: 100 (10 cassettes x 10 implants)

TO BE USED BY SUBCUTANEOUS EAR IMPLANTATION ONLY

Batch No.: Expiry Date:

> ADMINISTRATION: Approved implantation technique is fully described in the enclosed leaflet. WITHHOLDING PERIODS:

MEAT: NIL MILK: DO NOT use in heifers which are producing or may in the future produce milk or milk products

Scale: A3 = 77%

144x107x77 PSA020A04

L1195

x=10; y=50

123C 293C Black

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on 1800 226 511 before using this

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone 13 11 26*.

DO NOT dispose of unused implants in any way other than surrender to an approved

Dispose of empty cartridges by wrapping with paper and putting in garbage. Discarded needles should be placed in a designated 'sharps' container.

Store below 25°C (Air conditioning).

revalor[®] is recommended for use in non breeding cattle. It can be used during the growing or finishing phase and results in improved weight gain due to deposition of added lean meat. Repeat implantation can be carried out during the growing and finishing phase, however, it is recommended that repeat implantation is not carried out sooner than 70 days and that there is adequate nutrition to ensure a growth response. Growth response may vary widely for nutritional or other reasons. Good quality feed is a prerequisite for best results whether on pasture, supplement or lot feeding. Maximum growth rate requires good quality stock with adequate control of parasites and other diseases.





Scale: A4 = 100%

Dimensions: 13 x 45mm





CAUTION **KEEP OUT OF REACH OF CHILDREN** FOR ANIMAL TREATMENT ONLY

COOPERS®

Growth promotant for non breeding cattle

ACTIVE CONSTITUENTS:

EACH IMPLANT (4 PELLETS) CONTAINS: 80 mg TRENBOLONE ACETATE 16 mg 17 β OESTRADIOL

DIRECTIONS FOR USE LEAFLET:

To be read in conjunction with label affixed to outer package.

Restraints

FOR USE ONLY in non breeding cattle.

DO NOT use in heifers which are producing or may in the future produce milk or milk products for human consumption.

DO NOT USE this product in any other species of animal.

USE ONLY one implant per heifer or steer. Implant ONLY beneath the skin in the middle third of the ear.

DO NOT implant this product in any other site.

DO NOT USE this product on HGP free accredited cattle.

It is a breach of AQIS accreditation conditions to use this product on HGP free accredited cattle or to possess this product on any HGP free property (except for feedlots with flexible accreditation). Cattle implanted with this product must be individually identified with an approved ear punch mark.

DO NOT implant sooner than 70 days after last implant.

Implanting cattle under 6 weeks of age is NOT recommended.

Implanting cattle intended for breeding is NOT recommended as they may experience decreased fertility.

Growth responses may vary widely for nutritional or other reasons. Improved performance may be noted only when conditions are favourable for good growth.

Vaginal, rectal and preputial prolapse, increased bulling activity, high tailheads, sunken loins, ventral oedema and udder development may occur as side effects of treatment.

To help avoid abscess formation at the site of the implant, the surface of the ear should be thoroughly cleaned and disinfected before implantation.

MODE OF ACTION: Coopers REVALOR® FLEX is a combination of 2 growth promoting agents with differing modes of action. Trenbolone acetate is an anabolic steroid which is capable of increasing the accumulation of protein in animals by improving the nitrogen balance thus promoting the development of extra lean meat in the carcass.

 17β oestradiol is a naturally occurring hormone which also acts as a growth promoter but in a totally different manner via the brain and pituitary gland.

When used in combination in non breeding cattle, these compounds exert an additive effect, giving an improvement in growth rate and feed conversion superior to that of either component when used alone. The increase in bodyweight is predominantly lean meat and not fat.

This product is specifically designed and intended for non breeding cattle only. The implant is best administered 90 – 120 days before slaughter. It is especially effective under semi-intensive or intensive finishing conditions.

Scale: A4 = 100%

Dimensions: 125 x 210mm





DOSAGE: One implant is administered per animal in all cases. The 4 PELLETS which make up the implant of Coopers REVALOR® FLEX are contained in one division of the multiple cartridge. One implant is placed under the skin on the posterior aspect of the ear by means of the special Intervet implant gun.

ADMINISTRATION: The applicator:

Place the needle into position and tighten the fixing knurled nut, it is possible to orientate the bevelled edge of the needle into a convenient position with the bevel facing right for a right handed operator.



- a) Do not remove the cassette cover containing the implants.
- b) Insert the cassette into the upper aperture of the implant gun magazine (with the cover facing forwards).

In order to prevent damage to

g) Introduce the needle under

the skin, parallel with the

cartilage. If blood vessels

occurs, change location.

are damaged and bleeding

the implant, do ear tagging first.



- c) Press gently onto the cassette until a click is heard.
- d) The implant gun is ready for
- e) Ensure the skin on the outer surface of the ear is clean.
- f) Take hold of the animal's ear with your free hand (in the manner shown).



h) Press gently and progressively on the trigger so that the dose is injected (first part of the stroke) and the needle removed (second part of the stroke). The shield that rests on the skin at the implantation site causes the needle to retract automatically.



SITE OF IMPLANTATION

The administration should be made into the middle third of the back face of the ear, care being taken to avoid damaging large blood vessels or cartilage.



The sequence of the two operations is automatic and continuous and ensures that the pellets are deposited evenly in a straight line without risk of crushing or pilling up under the skin without being crushed in the needle.



- i) Remove the remaining tip of the needle and release the trigger.

i) The implant gun is then ready for the next animal.

When all implants have been administered, the cassette will fall out of the bottom of the gun magazine allowing replacement with a new cassette.

It is possible to remove a cassette that still contains several doses. It can be ejected by applying slight pressure movements successively on the trigger until the cassette clears the magazine.

CONCURRENT ADMINISTRATION OF OTHER VETERINARY DRUGS:

Other products e.g. anthelmintics, ecto-parasiticides, or antibiotics may be applied concurrently with Coopers REVALOR® FLEX.

Management practices such as ear tagging or branding should be carried out before implantation of Coopers REVALOR® FLEX to avoid damage to the implant.

WITHHOLDING PERIODS:

MEAT: NIL

MILK: DO NOT use in heifers which are producing or may in the future produce milk or milk products for human consumption.

EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For advice on the ESI, contact the manufacturer on 1800 226 511 before using this product.

If poisoning occurs, contact a doctor or the Poisons Information Centre. Phone 13 11 26.

DO NOT dispose of unused implants in any way other than surrender to an approved retailer/wholesaler.

Dispose of empty cartridges by wrapping with paper and putting in garbage. Discarded needles should be placed in a designated 'sharps' container.

STORAGE: Store below 25°C (Air conditioning).

For further information contact:

COOPERS ANIMAL HEALTH

A Division of Intervet Australia Pty Ltd

91-105 Harpin Street, BENDIGÓ EAST, VIC 3550 Phone: 1800 226 511

APVMA Approval No.: 58656/0610 revalor® is a Registered Trademark